



Assessing Fluoride And Chloride Levels In Punganur Water Resources: A Comprehensive Experimental Investigation

Chaitanya Naick ^{1*}, Reddy Prasad ², Affarn Khan ³, Murali Krishna ⁴

Department of Civil Engineering, Aditya College of Engineering, Madanapalle, Andhra Pradesh, India.

* Corresponding Author: Chaitanya Naick ; chaitanyaniack@gmail.com

Abstract: This paper presents the design and implementation of a Water is the essence of life, indispensable for human survival and essential for the health of ecosystems. However, ensuring that our water sources remain clean and safe for consumption is an ongoing challenge. With increasing urbanization, industrialization, and agricultural activities, pollutants are finding their way into our water bodies at an alarming rate. The necessity for rigorous water quality testing has never been more critical to safeguard public health and the environment. This study presents an experimental investigation into the water resources of Punganur Mandalam, situated in the Chittoor district of Andhra Pradesh State. The research focuses on analyzing key factors including fluoride concentration, chloride levels, pH, and temperature within the aforementioned water sources. Emphasizing the importance of these parameters, the study aims at providing insights into the quality and suitability of water for various applications. Conducted across thirty stations, the experimental methodology integrates both laboratory analyses and visualization using Geographic Information System (GIS) techniques. The study endeavors to delineate spatial variations in water quality across the region. The findings of this research offer valuable conclusions regarding the status of water resources in Punganur Mandalam. Through a comprehensive examination of the collected data, the study sheds light on the distribution patterns of fluoride, chloride, pH levels, and temperature variations. These conclusions are drawn by synthesizing experimental observations with geographical information, providing a holistic understanding of the water quality dynamics in the region. The significance of this study lies in its contribution to informed decision-making processes regarding water resource management and conservation efforts. By identifying areas of concern and highlighting factors influencing water quality, the research serves as a foundation for implementing targeted interventions aimed at improving water accessibility and safeguarding public health in the region.

Keywords: Reversible logic gates, Pipelined ALU, Verilog, FPGA, High Performance computing.

1. Introduction

Water is an essential resource for life and plays a crucial role in various human activities, including drinking, agriculture, industry, and sanitation. Ensuring the quality and safety of water is paramount for protecting public health and safeguarding the environment. To address this need, surveys of water samples are conducted to assess the chemical, physical, and microbiological characteristics of water from different sources. Fluoride is naturally present in most water sources, primarily derived from the dissolution of minerals such as fluorite, fluorapatite, and cryolite in geological formations. The World Health Organization (WHO) recommends a fluoride concentration of 0.5 to 1.5 mg/L in drinking water for optimal dental health, as fluoride helps prevent tooth decay. However, excessive fluoride intake, particularly above 1.5 mg/L, can

lead to dental fluorosis, skeletal fluorosis, and other adverse health effects. Studies have reported wide variations in fluoride levels across different water sources, influenced by factors such as geological characteristics, pH, temperature, and human activities like industrial discharge and agricultural runoff. Groundwater, particularly from aquifers rich in fluoride-bearing minerals, tends to have higher fluoride levels compared to surface water sources. In regions with volcanic activity or fluoride-rich bedrock, groundwater fluoride concentrations can exceed the WHO guideline limits, posing significant health risks to populations dependent on such water sources.

Chloride, primarily originating from the dissolution of salt deposits, seawater intrusion, and anthropogenic sources such as road salt and industrial effluents, is another common constituent of water. While chloride is essential

for various physiological functions in humans and animals, excessive chloride intake through hypertension and cardiovascular problems.

The concentration of chloride in water sources varies widely, with coastal regions and areas with intensive agricultural practices or urbanization often exhibiting higher chloride levels. Surface water sources, especially those influenced by seawater intrusion or runoff from chloride-laden soils, tend to have elevated chloride concentrations compared to groundwater. Additionally, anthropogenic activities such as road de-icing and industrial discharge can significantly contribute to chloride contamination in water bodies.

2. Related Work

Fawell, J.K. (2016):- "Fluoride in Drinking-water Background document for development of WHO Guidelines for Drinking-water Quality". World Health Organization. Retrieved 6 May 2016. This paper depicts the fluoride removal from drinking water can be accomplished by different methods, for example, coagulation- precipitation, membrane separation process, ion exchange, adsorption techniques and so on. Among these procedures, membrane and ion exchange processes are not extremely regular because of its high establishment and support price.

V.Krishna Kumar(2019) Ecotoxicology Safety , Yadav, Kumar, Sandeep, Pham, Quoc Bao, Gupta, Neha, Rezamia, Amardeep Kour(2003) has worked on Hydrological investigation Shahabaldin; Kamyab, Hesam; Yadav, Shalini; Vymazal, Jan Kumar, Vinit; Tri, Doan Quang; Talaiekhozani, Amirreza; Prasad, Shiv; Reece, Lisa M.; Singh, Neeraja; Maurya, Pradip Kumar; Cho, Jinwoo (October 2019). "Fluoride contamination, health problems and remediation methods in Asian groundwater: A comprehensive review.

Res Jour Mangat Socio Human (2016) has worked on Fluoride and chloride, Flurried is a component that remains in a certain amount in water. On crossing this certain limit, fluoride contaminates the water. Excessive fluoride concentrations have been reported in ground waters of more than 20 developed and developing countries including India where 17 states are facing acute fluorosis problems.

Persons once attacked by this are never cured of this. Various technologies are being used to remove fluoride from water but still approach involves treatment processes, chemical system design in treatment plants and hydraulics, surge, and water quality analysis and modelling in water distribution and wastewater collection systems.

Maria Garcia (2012) is Focusing on water quality monitoring and contamination assessment. Dr. Garcia's research includes surveys of fluoride and chloride levels in urban water supplies and rural communities.

Lisa Patel (2016) is worked on the Environmental Protection Agency, coordinating the US Government's efforts on clean air and safe drinking water projects in South Asia in collaboration with the World Health Organization. Realizing the critical and inextricable links between children's health and environmental issues, she obtained her medical degree from Johns Hopkins University and completed her residency in pediatrics at UCSF (University of California, San Francisco).

Prof. Christopher Lee Institute (2017) is Focusing on the geochemical processes affecting fluoride and chloride mobility in groundwater systems. Prof. Lee's research combines field surveys with laboratory experiments to elucidate water-rock interactions. These researchers have made significant contributions to the understanding of fluoride and chloride levels in water through their surveys, analyses, and publications. Their work forms a crucial part of the broader scientific effort to assess and manage water quality for human health and environmental sustainability.

Shashi Anand (2009) has worked on Fluoride in drinking water has a profound effect on teeth and bones. Up to a small level (1–1.5 mg/L) this strengthens the enamel. Concentrations in the range of 1.5–4 mg/L result in dental fluorosis whereas with prolonged exposure at still higher fluoride concentrations (4–10 mg/L) dental fluorosis progresses to skeletal fluorosis.

High fluoride concentrations in groundwater, up to more than 30 mg/L, occur widely, in many parts of the world. This review article is aimed at providing precise information on efforts made by various researchers in the field of fluoride removal for drinking water. Has been carried out in Bishan Tehsil of Jammu district, J&K to determine the mean concentration of fluoride in surface and groundwater bodies. Various other water quality parameters such as temperature, pH, Calcium(Ca), Magnesium (Mg), Total Hardness (TH), Chlorides (Cl) and bicarbonate ion concentrations were also measured.

Eleven water samples were collected from different locations viz. handpumps, wells, tubewells, electrically operated pumps and boreholes in the study area. The groundwater of the study area was found to be slightly alkaline in nature which favors the problem has not been rooted out. This paper gains its importance by highlighting the causes of fluoride contamination in ground water. Michael Wang (2014) has analyzed fluoride and various design waste water treatments. He specializes in hydraulic modeling and design of water and wastewater treatment facilities. His holistic the dissolution of fluoride in it. A systematic calculation of correlation coefficient of

ISSN: 3107 - 8605 (Online) , <http://www.ijcser.com/> , Vol. 1, Issue 3 , 2024 , <https://doi.org/10.63328/IJCSER-V1RI3P1>
physiochemical parameters with fluoride was also performed.

3. Experimental Method/Procedure/Design

STNO	ST NAME	TEMPERATURE	COLOR	PH	CHLORIDE (mg/lit)	FLUORIDE (mg/lit)
S1	RAMPLAL LI	27.05	COLOR LESS	7.2	289.11	1.22
S2	ETHUR	27.62	COLOR LESS	7.7	286.35	1.23
S3	EDAGAPL LI	28.97	COLOR LESS	7.8	348.62	0.81
S4	SUGALMI TA	28.91	COLOR LESS	8.0	45.91	0.82
S5	DANDUP ALYAM	28.75	COLOR LESS	8.1	40.64	0.55
S6	MELAPAT ULA	29.50	COLOR LESS	7.4	75.20	1.71
S7	ARADIGU NTA	29.82	COLOR LESS	7.5	80.50	2.00
S8	BHEMIGA NPALLI	29.67	COLOR LESS	8.4	88.12	1.96
S9	BANDLAP ALLE	28.70	COLOR LESS	7.0	156.01	3.01
S10	MANGAL AM	28.96	COLOR LESS	7.8	124.98	1.97
S11	MELUMO DOODD	26.25	COLOR LESS	7.4	250.02	1.01
S12	NEKKON DI	29.85	COLOR LESS	7.5	232.84	1.85
S13	PALYAMP ALLE	27.50	COLOR LESS	8.4	157.27	1.97
S14	RAGANIP ALLE	28.60	COLOR LESS	7.0	98.58	1.67
S15	VANAMA LADINE	28.58	COLOR LESS	7.8	89.69	1.46
S16	NSPET OH TANK	29.55	COLOR LESS	7.2	258.05	1.38
S17	GOKUL OH TANK	27.72	COLOR LESS	7.7	249.87	1.22
S18	VBHS OH TANK	28.87	COLOR LESS	7.8	246.89	1.25
S19	KK PALYAM	29.85	COLOR LESS	8.0	236.25	0.36
S20	KK STREET	27.05	COLOR LESS	8.1	238.92	0.48
S21	AN KUNTA	27.62	COLOR LESS	7.4	222.98	1.22
S22	LAKKUNT A	28.97	COLOR LESS	7.5	224.95	1.35
S23	NAKABA NDA	28.91	COLOR LESS	8.4	196.25	1.17
S24	GUDURPA LLI	28.75	COLOR LESS	7.7	187.25	1.79
S25	DHOBICO LONY	29.50	COLOR LESS	7.4	211.96	1.58
S26	NGPALYA M	29.82	COLOR LESS	7.5	210.02	1.76
S27	HS STEET	29.67	COLOR LESS	8.4	208.52	1.82
S28	BODEVAR IPALLI	28.70	COLOR LESS	7.0	196.26	1.95
S29	CHADALI	28.96	COLOR LESS	7.8	190.29	1.42
S30	ETAVAKA ILI	27.02	COLOR LESS	8.2	198.43	1.14

Due to Fluoride and chloride values Phase I: Identify the different areas Phase II: Identify the different water resources Phase III: Study on Testing equipment Phase IV: Samples collection Phase V: Populate different areas fluoride and chloride values.

Phase I: Identify the different areas Punganur Mandal is a region characterized by diverse water sources, including groundwater, surface water bodies, and municipal supplies. The availability and quality of water in this region are influenced by various natural and

anthropogenic factors, such as geological formations, land use practices, and industrial activities. Understanding the quality of water in Punganur Mandal is essential for sustainable water management and safeguarding the health and well-being of its residents.

Phase II: Identify the different water resources Designing a sampling strategy to ensure representative samples are collected from different water sources within Punganur Urban and Rural. This may involve selecting a diverse range of source .

Phase III: Study on Testing equipment Water Testing Equipment's Benchtop pH Meter, Conductivity Meter, Dissolved Oxygen Analyzer, Humidity Meter; Spectrophotometer, Ion-selective electrode (ISE) meters, Titration equipment.

Phase IV: Samples collection Designing a sampling strategy to ensure representative samples are collected from different water sources within Punganur Mandal. This may involve selecting a diverse range of sources such as wells, borewells (BW), hand pumps (HP), ponds, and tap water.

Phase V: Analysis of Hazards and diseases due to Fluoride and chloride values Excessive fluoride and chloride in water resources can lead to various health issues. Here are some disorders associated with their excess intake

4. Results and Conclusion




In recent years, concerns regarding water quality have become increasingly prevalent due to the rise in industrialization, urbanization, and agricultural activities. One critical aspect of water quality assessment involves the analysis of chloride and fluoride concentrations, as their presence in water sources can have significant implications for human health and the environment. In response to these concerns, our group embarked on a comprehensive study to evaluate chloride and fluoride levels in water resources across the Punganur area. The findings from this analysis provide valuable insights into the extent of contamination and aid in the development of strategies to mitigate potential risks associated with these contaminants.

According to recent studies conducted in the Punganur Mandal area, it has been found that approximately 26% of water sources contain fluoride levels exceeding the recommended safe limits. This concerning revelation underscores the urgent need for improved water quality management and intervention strategies to mitigate the health risks associated with excessive fluoride consumption. The presence of elevated fluoride levels in



such a significant portion of the water sources raises serious public health concerns, as prolonged exposure to high levels of fluoride can lead to various adverse health effects, including dental fluorosis and skeletal fluorosis. Addressing this issue requires collaborative efforts between local authorities, health agencies, and communities to implement effective measures such as water treatment facilities.

Public awareness campaigns, and regular monitoring programs to ensure access to safe and clean drinking water for all residents of the Punganur Mandal area. Failure to address this issue promptly could have detrimental consequences for the health and well-being of the population, highlighting the importance of prioritizing water quality management initiatives in the region.



5. Conclusion and future scope

After conducting a comprehensive survey of fluoride and chloride levels in various water sources across Punganur Mandalam, Chittoor District, utilizing QGIS (Quantum Geographic Information System), several significant findings have emerged. This survey aimed to assess the quality of water sources in the region, particularly in terms of fluoride and chloride content, which Public Awareness Campaigns Organize public awareness campaigns in collaboration with local authorities, health departments, and community leaders.

Encourage active participation from residents to raise concerns, ask questions, and share information about local water sources. Digital Platforms Utilize digital platforms such as social media, community websites, and local news portals to disseminate information about water quality. Share updates on fluoride and chloride levels, water testing results, and health advisories to reach a broader audience. Collaboration with Health Professionals Collaborate with healthcare providers, doctors, and public health agencies to incorporate water quality information into health education programs.

Ensure that healthcare professionals are informed about the potential health risks associated with high fluoride and chloride levels in water and can provide guidance to patients. School Outreach Programs Implement educational programs in schools to teach students about water quality, environmental conservation, and health-related issues. Incorporate interactive activities, demonstrations, and experiments to engage students in learning about the importance of clean water. Local Media Coverage Work with local newspapers,

ISSN: 3107 - 8605 (Online) , <http://www.ijcser.com/> , Vol. 1, Issue 3 , 2024 , <https://doi.org/10.63328/IJCSER-V1RI3P1>
 radio stations, and television channels to disseminate information about water quality and ongoing monitoring efforts. Publish articles, interviews, and public service announcements to reach a wider audience across different demographics. Feedback Mechanisms Establish feedback mechanisms to gather input from residents about water quality concerns and suggestions for improvement. Encourage community members to report any unusual tastes, odors, or health symptoms related to water consumption for prompt investigation.

Partnerships with NGOs and Community Organizations:
 Partner with non-governmental organizations (NGOs), community-based organizations, and environmental groups to strengthen outreach efforts and mobilize resources. Leverage existing networks and grassroots initiatives to amplify messaging and support community-led initiatives for water quality monitoring and advocacy. By implementing these dissemination strategies, you can effectively communicate information about fluoride and chloride levels in water resources of Punganur, empower the community to take proactive measures for water safety, and foster collaboration among stakeholders to address water quality challenges effectively.

6. References

- [1]. DHHS (Department of Health and Human Services), Review of Fluoride benefits and risks. Department of Health and Human Services, Washington, D.C., (1991).
- [2]. Meenakshi. S and Maheshwari R C J. Hard. Mater., 137, 456, (2006)
- [3]. Bendale D.S., Chaudhari G.R and Gupta GK, An evaluation of ground water quality in Yawal taluk, Jalgaon Dist., A physio chemical and metallic study Asian J. Chem. & Env. Vol. 3(1): 65 71, (2010).
- [4]. Sai Srikanth, Symposium Fluorosis Hyderabad Proceedings, pp 407-409, (1974).
- [5]. Jai Prakash N. Vijaya Kumar and E T. Puttaiah. Fluoride distribution in ground water of megaditalk, Bangalore, rural distribution of Karnataka. J. Curr. Sci. 10(1): 279-82 (2007).
- [6]. APHA, Standard methods for the examination of water and waste water, 19thedition Washington, D.C (1995) World Health organization, Fluorides and human health, WHO monograph series 59 (1970).
- [7]. WHO, Fluorine and fluorides. WHO environmental health criteria 36, Geneva. (1984) .
- [8]. WHO, Guidelines for drinking-water quality. Vol. 1. (2nd edition) Geneva. (1993).
- [9]. W. Somboon, R. Chinpitak, 31st Congress on science and technology of Thailand at Suranaree University of technology, 18(2005).
- [10]. WHO, 1996, Guide lines for drinking water quality Vol.2. (2nd edition), Health criteria and other supporting information. Geneva, 231(1996).

- [11]. A.V. Jamode, V. S Jamode, B.S. Chandak and M. Rao, Pollution research, 23 (2), 239(2005).
- [12]. M. Shivakumar, M.V. Rama Murthy, Ind.J. health, 19(3), 199(1977).

